

About amenability of subgroups of the group of diffeomorphisms of the interval.

E.T.Shavgulidze

*Department of Mechanics and Mathematics, Moscow State University, Moscow,
119991 Russia*

Averaging linear functional on the space continuous functions of the group of diffeomorphisms of interval is found. Amenability of several discrete subgroups of the group of diffeomorphisms $\text{Diff}^3([0, 1])$ of interval is prove. In particular, a solution of the problem of amenability of the Thompson's group F is given.

1. The main result.

Let $\text{Diff}_+^1([0, 1])$ be the group of all diffeomorphisms of class C^1 of interval $[0, 1]$ that preserve the endpoints of interval, and let $\text{Diff}_+^3([0, 1])$ be the subgroup of $\text{Diff}_+^1([0, 1])$ consisting of all diffeomorphisms of class $C^3([0, 1])$, and

$$\text{Diff}_0^3([0, 1]) = \{f \in \text{Diff}_+^3([0, 1]) : f'(0) = f'(1) = 1\}.$$

The group $\text{Diff}_+^1([0, 1])$ is equipped with the topology inherited from the space $C^1([0, 1])$.

Let us denote $D_n = \{(x_1, \dots, x_{n-1}) : 0 < x_1 < \dots < x_{n-1} < 1\} \subset \mathbf{R}^{n-1}$ and $x_0 = 0, x_n = 1$.

We say that a subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a), if

(i) there are a integer $l_n > l_{n-1}$ ($l_0 = 1$) for any natural n and a countably additive Borel measure η_n on D_{l_n} such that $\eta_n(D_{l_n}) = 1$,

(ii) for any positive ε and any $g \in G$, we can find natural $N(\varepsilon, g)$ such that, for any $n > N(\varepsilon, g)$, it exists a Borel subset $Z_{n,\varepsilon,g} \subset D_{l_n}$ that $\eta_n(Z_{n,\varepsilon,g}) > 1 - \varepsilon$ and $\max_{1 \leq k \leq l_n} (x_k - x_{k-1}) < \varepsilon$ for any $(x_1, x_2, \dots, x_{l_n-1}) \in Z_{n,\varepsilon,g}$ where $x_0 = 0, x_{l_n} = 1$,

(iii) $(1 - \varepsilon)\eta_n(Y) < \eta_n(gY) < (1 + \varepsilon)\eta_n(Y)$ for any Borel subset $Y \subset Z_{n,\varepsilon,g}$ where $gY = \{(g(x_1), g(x_2), \dots, g(x_{l_n-1})) : (x_1, x_2, \dots, x_{l_n-1}) \in Y\}$.

For any positive $\delta < 1$, denote by $C_0^{1,\delta}([0, 1])$ the set of all functions $f \in C^1([0, 1])$ such that $f(0) = 0$ and $\exists C > 0 \forall t_1, t_2 \in [0, 1] |f'(t_2) - f'(t_1)| < C|t_2 - t_1|^\delta$. Define a Banach structure on the linear space $C_0^{1,\delta}([0, 1])$ by a norm

$$\|f\|_{1,\delta} = |f'(0)| + \sup_{t_1, t_2 \in [0, 1]} \frac{|f'(t_2) - f'(t_1)|}{|t_2 - t_1|^\delta}$$

for any function $f \in C_0^{1,\delta}([0, 1])$.

Let $\text{Diff}_+^{1,\delta}([0, 1]) = \text{Diff}_+^1([0, 1]) \cap C_0^{1,\delta}([0, 1])$. It is easy to see that $\text{Diff}_+^{1,\delta}([0, 1])$ is a subgroup of the group $\text{Diff}_+^1([0, 1])$. The subgroup $\text{Diff}_+^{1,\delta}([0, 1])$ is equipped with the topology inherited from the space $C_0^{1,\delta}([0, 1])$.

Let $C_b(\text{Diff}_+^{1,\delta}([0, 1]))$ be the linear space of all bounded continuous functions on the space $\text{Diff}_+^{1,\delta}([0, 1])$, and let $C_b(\text{Diff}_+^1([0, 1]))$ be the linear space of all bounded continuous functions on the space $\text{Diff}_+^1([0, 1])$.

Introduce the functions $e_{1,\delta} : \text{Diff}_+^{1,\delta}([0, 1]) \rightarrow \mathbf{R}$, $e_{1,0} : \text{Diff}_+^1([0, 1]) \rightarrow \mathbf{R}$ by setting $e_{1,\delta}(g) = 1$ for any $g \in \text{Diff}_+^{1,\delta}([0, 1])$ and $e_{1,0}(f) = 1$ for any $f \in \text{Diff}_+^1([0, 1])$. Let $F_g(f) = F(g^{-1} \circ f)$ for any $g \in \text{Diff}_0^3([0, 1])$, $f \in \text{Diff}_+^{1,\delta}([0, 1])$ and $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$.

Theorem 1. *If a subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a) and a positive $\delta < \frac{1}{2}$ then there exists a linear functional*

$$L_\delta : C_b(\text{Diff}_+^{1,\delta}([0, 1])) \rightarrow \mathbf{R} \text{ such that } L_\delta(e_{1,\delta}) = 1, |L_\delta(F)| \leq \sup_{f \in \text{Diff}_+^{1,\delta}([0, 1])} |F(f)|,$$

$L_\delta(F) \geq 0$ for any nonnegative function $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$, and

$$L_\delta(F_g) = L_\delta(F) \text{ for any } g \in G \text{ and } F \in C_b(\text{Diff}_+^{1,\delta}([0, 1])).$$

The restriction of any function of the space $C_b(\text{Diff}_+^1([0, 1]))$ on $\text{Diff}_+^{1,\delta}([0, 1])$ belongs to the space $C_b(\text{Diff}_+^{1,\delta}([0, 1]))$. Hence we obtain the following assertion.

Corollary 1.1. *If a subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a) then there exists a linear functional $L_0 : C_b(\text{Diff}_+^1([0, 1])) \rightarrow \mathbf{R}$ such that $L_0(e_{1,0}) = 1$, $|L_0(F)| \leq \sup_{f \in \text{Diff}_+^1([0, 1])} |F(f)|$, $L_0(F) \geq 0$ for any nonnegative function $F \in C_b(\text{Diff}_+^1([0, 1]))$, and $L_0(F_g) = L_0(F)$ for any $g \in G$ and $F \in C_b(\text{Diff}_+^1([0, 1]))$.*

We say that a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (b), if there is a such $C > 0$ that

$$\sup_{t \in [0, 1]} |\ln(g'_1(t)) - \ln(g'_2(t))| \geq C \text{ for any } g_1, g_2 \in G, g_1 \neq g_2.$$

Theorem 2. *If a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies conditions (a), (b), then the subgroup G is amenable.*

In [2] È.Ghys and V.Sergiescu proved that the Thompson's group F is isomorphic to a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ which satisfies condition (b).

Corollary 2.1. *The Thompson's group F is amenable.*

2. Proof of Theorem 1.

Define the mapping $A : \text{Diff}_+^1([0, 1]) \rightarrow C_0([0, 1])$ by setting

$$A(q)(t) = \ln(q'(t)) - \ln(q'(0)) \quad \forall t \in [0, 1].$$

The mapping A is a topological isomorphism between the space $\text{Diff}_+^1([0, 1])$, $C_0([0, 1])$ moreover

$$A^{-1}(\xi)(t) = \frac{\int_0^t e^{\xi(\tau)} d\tau}{\int_0^1 e^{\xi(\tau)} d\tau}.$$

Introduce the Wiener measure w on the space $C_0([0, 1])$. Define a Borel measure ν on $\text{Diff}_+^1([0, 1])$ by setting $\nu(X) = w(A(X))$ for any Borel subset X of topological space $\text{Diff}_+^1([0, 1])$.

Let $\delta \in (0, \frac{1}{2})$. It follows from the properties of Wiener measure w (see [4]) that measure ν is concentrated on the set $E_\delta = \text{Diff}_+^{1,\delta}([0, 1])$, i.d. $\nu(E_\delta) = 1$, moreover the Borel subsets of metric space E_δ is measurable with respect to the measure ν .

As it was proved in [3], the measure ν is quasi-invariant with respect to the left action of subgroup $\text{Diff}_+^3([0, 1])$ on the group $\text{Diff}_+^1([0, 1])$, moreover

$$\nu(gX) = \frac{1}{\sqrt{g'(0)g'(1)}} \int_X e^{\frac{g''(0)}{g'(0)}q'(0) - \frac{g''(1)}{g'(1)}q'(1) + \int_0^1 S_g(q(t))(q'(t))^2 dt} \nu(dq),$$

for any Borel subset X of topological space $\text{Diff}_+^1([0, 1])$, and any $g \in \text{Diff}_+^3([0, 1])$, where $gX = \{g \circ q : q \in X\}$ and $S_g(\tau) = \frac{g'''(\tau)}{g'(\tau)} - \frac{3}{2}(\frac{g''(\tau)}{g'(\tau)})^2$ (the Schwartz derivative of function g).

For the proof of Theorem 1 we need the following auxiliary assertions

Lemma 1. *The following equality is valid*

$$\int_{E_\delta} (q'(0))^l \nu(dq) = \int_{E_\delta} (q'(1))^l \nu(dq)$$

for any natural l .

Proof. Let $\xi = A(q)$, i.d. $\xi(t) = \ln(q'(t)) - \ln(q'(0))$. Then

$$q'(0) = \frac{1}{\int_0^1 e^{\xi(\tau)} d\tau}, \quad q'(1) = \frac{e^{\xi(1)}}{\int_0^1 e^{\xi(\tau)} d\tau}.$$

Let us take

$$\begin{aligned} M_l &= \int_{E_\delta} (q'(1))^l \nu(dq) = \int_{\text{Diff}_+^1([0,1])} (q'(1))^l \nu(dq) = \\ &= \int_{C_0([0,1])} \left(\frac{e^{\xi(1)}}{\int_0^1 e^{\xi(\tau)} d\tau} \right)^l w(d\xi) = \int_{C_0([0,1])} \left(\frac{1}{\int_0^1 e^{\xi(1-\tau)-\xi(1)} d\tau} \right)^l w(d\xi) \end{aligned}$$

Let $\zeta(t) = \xi(1-t) - \xi(1)$. The Wiener measure w is invariant with respect to the action $\zeta \mapsto \xi$, thus,

$$\begin{aligned} M_l &= \int_{C_0([0,1])} \left(\frac{1}{\int_0^1 e^{\zeta(\tau)} d\tau} \right)^l w(d\zeta) = \\ &= \int_{\text{Diff}_+^1([0,1])} (q'(0))^l \nu(dq) = \int_{E_\delta} (q'(0))^l \nu(dq), \end{aligned}$$

which implies the assertion of Lemma 1.

Introduce the measure $\nu_n = \nu \otimes \dots \otimes \nu$ on the space $E_\delta^n = E_\delta \times \dots \times E_\delta$.

Let $c_1 = 1 + M_1 + M_2 + \int_{E_\delta} (\int_0^1 (q'(t))^2 dt) \nu(dq)$.

For any $r > 0$, $g \in \text{Diff}_+^3([0,1])$, $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$, we write $C_g = 1 + \max_{0 \leq t \leq 1} (|\frac{g''(t)}{g'(t)}| + (\frac{g''(t)}{g'(t)})^2 + |\frac{g'''(t)}{g'(t)}|)$ and

$$\begin{aligned} X_{r,g,\bar{x}} &= \{(q_1, \dots, q_n) : q_1, \dots, q_n \in E_\delta \\ &| \sum_{k=1}^n [(x_k - x_{k-1}) (\frac{g''(x_{k-1})}{g'(x_{k-1})} q'_k(0) - \frac{g''(x_k)}{g'(x_k)} q'_k(1)) + \\ &+ (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1}) q_k(t)) (q'_k(t))^2 dt] \leq 4c_1 C_g r\}. \end{aligned}$$

Lemma 2. *If $\epsilon \in (0,1)$, then the following inequality is fulfilled*

$\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon}, g, \bar{x}}) \leq 2\sqrt[3]{\epsilon}$ for any $g \in \text{Diff}_+^3([0,1])$, for any positive integer n and $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$, satisfying the inequality $\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \epsilon$.

Proof. Let

$$f_1(q_1, \dots, q_n) = \sum_{k=1}^n (x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} q'_k(0) - \frac{g''(x_k)}{g'(x_k)} q'_k(1) \right).$$

Then

$$I_1 = \int_{E_\delta} \dots \int_{E_\delta} f_1(q_1, \dots, q_n) \nu(dq_1) \dots \nu(dq_n) = M_1 \sum_{k=1}^n (x_k - x_{k-1}) \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} - \frac{g''(x_k)}{g'(x_k)} \right).$$

As $\left| \frac{g''(x_{k-1})}{g'(x_{k-1})} - \frac{g''(x_k)}{g'(x_k)} \right| \leq C_g (x_k - x_{k-1})$, we have

$$|I_1| \leq M_1 C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \leq M_1 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = M_1 C_g \epsilon.$$

If $k \neq l$ then

$$\begin{aligned} & \int_{E_\delta} \int_{E_\delta} \left(\frac{g''(x_{k-1})}{g'(x_{k-1})} (q'_k(0) - M_1) - \frac{g''(x_k)}{g'(x_k)} (q'_k(1) - M_1) \right) \\ & \left(\frac{g''(x_{l-1})}{g'(x_{l-1})} (q'_l(0) - M_1) - \frac{g''(x_l)}{g'(x_l)} (q'_l(1) - M_1) \right) \nu(dq_k) \nu(dq_l) = 0, \end{aligned}$$

therefore

$$\begin{aligned} I_2 &= \int_{E_\delta} \dots \int_{E_\delta} (f_1(q_1, \dots, q_n) - I_1)^2 \nu(dq_1) \dots \nu(dq_n) = \\ & \sum_{k=1}^n (x_k - x_{k-1})^2 \int_{E_\delta} \left[\frac{g''(x_{k-1})}{g'(x_{k-1})} (q'_k(0) - M_1) - \frac{g''(x_k)}{g'(x_k)} (q'_k(1) - M_1) \right]^2 \nu(dq_k) \leq \\ & \leq 2 \sum_{k=1}^n (x_k - x_{k-1})^2 \left[\left(\frac{g''(x_{k-1})}{g'(x_{k-1})} \right)^2 \int_{E_\delta} (q'_k(0) - M_1)^2 \nu(dq_k) + \right. \\ & \quad \left. + \left(\frac{g''(x_k)}{g'(x_k)} \right)^2 \int_{E_\delta} (q'_k(1) - M_1)^2 \nu(dq_k) \right] = \\ & = 2 \sum_{k=1}^n (x_k - x_{k-1})^2 \left[\left(\frac{g''(x_{k-1})}{g'(x_{k-1})} \right)^2 + \left(\frac{g''(x_k)}{g'(x_k)} \right)^2 \right] \left[\int_{E_\delta} (q'_k(0))^2 \nu(dq_k) - (M_1)^2 \right] \leq \\ & \leq 4M_2 C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \leq 4M_2 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = 4M_2 C_g \epsilon. \end{aligned}$$

Hence,

$$\nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n) - I_1| \geq 2c_4 C_g \sqrt[3]{\epsilon}\}) \leq \frac{I_2}{(2c_4 C_g \sqrt[3]{\epsilon})^2} \leq \frac{4M_2 C_g \epsilon}{(2c_4 C_g \sqrt[3]{\epsilon})^2} \leq \sqrt[3]{\epsilon}.$$

Thus

$$\nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n)| \geq 3c_1 C_g \sqrt[3]{\epsilon}\}) \leq \sqrt[3]{\epsilon}.$$

Let

$$f_2(q_1, \dots, q_n) = \sum_{k=1}^n ((x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t))(q'_k(t))^2 dt.$$

Then

$$I_3 = \int_{E_\delta} \dots \int_{E_\delta} |f_2(q_1, \dots, q_n)| \nu(dq_1) \dots \nu(dq_n) \leq$$

$$\begin{aligned}
&\leq 2C_g \sum_{k=1}^n (x_k - x_{k-1})^2 \int_{E_\delta} \left(\int_0^1 (q'_k(t))^2 dt \right) \nu(dq_k) \leq \\
&\leq 2c_1 C_g \epsilon \sum_{k=1}^n (x_k - x_{k-1}) = 2c_1 C_g \epsilon.
\end{aligned}$$

Thus

$$\nu_n(\{(q_1, \dots, q_n) : |f_2(q_1, \dots, q_n)| \geq 2c_1 C_g \sqrt[3]{\epsilon}\}) \leq \frac{I_3}{2c_1 C_g \sqrt[3]{\epsilon}} \leq \frac{2c_1 C_g \epsilon}{2c_1 C_g \sqrt[3]{\epsilon}} = (\sqrt[3]{\epsilon})^2 \leq \sqrt[3]{\epsilon}.$$

Hence,

$$\begin{aligned}
&\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon}, g, \bar{x}}) = \\
&= \nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n) + f_2(q_1, \dots, q_n)| \geq 4c_1 C_g \sqrt[3]{\epsilon}\}) \leq \\
&\leq \nu_n(\{(q_1, \dots, q_n) : |f_1(q_1, \dots, q_n)| \geq 2c_1 C_g \sqrt[3]{\epsilon}\}) + \\
&+ \nu_n(\{(q_1, \dots, q_n) : |f_2(q_1, \dots, q_n)| \geq 2c_1 C_g \sqrt[3]{\epsilon}\}) \leq 2\sqrt[3]{\epsilon},
\end{aligned}$$

which implies the assertion of Lemma 2.

Lemma 3. For any $g \in \text{Diff}_0^3([0, 1])$, $\epsilon > 0$, there is $\delta_1 \in (0, 1)$ such that the inequality is valid

$$\left| \prod_{k=1}^n \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1}) \sqrt{g'(x_k)g'(x_{k-1})}} - 1 \right| \leq \epsilon$$

for any natural n and any $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequality

$$\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \delta_1, \text{ where } x_0 = 0, x_n = 1.$$

Proof. Let $\epsilon \in (0, 1)$. Let $C = \max_{t_1, t_2 \in [0, 1]} (1 + |\frac{g''(t_1)}{g'(t_1)}| + |\frac{g'''(t_2)}{g'(t_1)}|)^2$, $\delta_1 = \frac{1}{400(C+1)}$,

$x'_k = \frac{x_k - x_{k-1}}{2}$ for any k ($1 \leq k \leq n$).

There are $x_k^*, x_k^{**}, x_k^{***} \in (0, 1)$ such that

$$\begin{aligned}
g(x_k) - g(x_{k-1}) &= g'(x'_k)(x_k - x_{k-1}) + \frac{1}{24} g'''(x_k^*)(x_k - x_{k-1})^3 = \\
&= g'(x'_k)(x_k - x_{k-1}) \left(1 + \frac{g'''(x_{k-1}^*)}{24g'(x'_k)} (x_k - x_{k-1})^2 \right), \\
g'(x_k) &= g'(x'_k) + \frac{1}{2} g''(x'_k)(x_k - x_{k-1}) + \frac{1}{8} g'''(x_k^{**})(x_k - x_{k-1})^2 \\
&= g'(x'_k) \left(1 + \frac{g''(x'_k)}{2g'(x'_k)} (x_{k-1} - x_{k-2}) + \frac{g'''(x_k^{**})}{8g'(x'_k)} (x_{k-1} - x_{k-2})^2 \right), \\
g'(x_{k-1}) &= g'(x'_k) - \frac{1}{2} g''(x'_k)(x_k - x_{k-1}) + \frac{1}{8} g'''(x_k^{***})(x_k - x_{k-1})^2 \\
&= g'(x'_k) \left(1 - \frac{g''(x'_k)}{2g'(x'_k)} (x_{k-1} - x_{k-2}) + \frac{g'''(x_k^{***})}{8g'(x'_k)} (x_{k-1} - x_{k-2})^2 \right).
\end{aligned}$$

Hence

$$\frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1}) \sqrt{g'(x_k)g'(x_{k-1})}} = \frac{1 + \lambda'_k(x_k - x_{k-1})}{\sqrt{1 + \lambda''_k(x_k - x_{k-1})}}$$

where

$$\begin{aligned}
\lambda'_k &= \frac{g'''(x_{k-1}^*)}{24g'(x'_k)} (x_k - x_{k-1}), \\
\lambda''_k &= \left(\frac{g''(x'_k)}{2g'(x'_k)} \right)^2 + \frac{g'''(x_k^{**}) + g'''(x_k^{***})}{8g'(x'_k)} (x_k - x_{k-1}) +
\end{aligned}$$

$$+ \frac{g''(x'_k)(g'''(x_k^{***}) - g'''(x_k^{**}))}{16(g'(x'_k))^2}(x_k - x_{k-1})^2 + \frac{g'''(x_k^{***})g'''(x_k^{**})}{64(g'(x'_k))^2}(x_k - x_{k-1})^3.$$

As $(x_k - x_{k-1}) < \delta_1$, there are $|\lambda'_k| < C\delta_1 < \frac{\epsilon}{100}$, $|\lambda''_k| < C\delta_1 < \frac{\epsilon}{100}$.

We have

$$\begin{aligned} \sigma &= \ln\left(\prod_{k=1}^n \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1})\sqrt{g'(x_k)g'(x_{k-1})}}\right) = \\ &= \sum_{k=1}^n (\ln(1 + \lambda'_k(x_k - x_{k-1})) - \frac{1}{2} \ln(1 + \lambda''_k(x_k - x_{k-1}))) \end{aligned}$$

and

$$|\sigma| \leq 2 \sum_{k=1}^n (|\lambda'_k| + |\lambda''_k|)(x_k - x_{k-1}) \leq \frac{\epsilon}{10} \sum_{k=1}^n (x_k - x_{k-1}) = \frac{\epsilon}{10},$$

therefore

$$\left| \prod_{k=1}^n \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1})\sqrt{g'(x_k)g'(x_{k-1})}} - 1 \right| = |e^\sigma - 1| \leq e^{\frac{\epsilon}{10}} - e^{-\frac{\epsilon}{10}} \leq \frac{\epsilon}{5} + \frac{\epsilon}{5} < \epsilon,$$

which implies the assertion of Lemma 3.

Introduce the mapping $Q_n : D_n \times E_\delta^n \rightarrow E_\delta = \text{Diff}_+^{1,\delta}([0, 1])$ by setting $f_n \circ (\tilde{l}_n)^{-1} = Q_n(x_1, \dots, x_{n-1}, \varphi_1, \dots, \varphi_n)$, where

$$f_n(t) = x_{k-1} + (x_k - x_{k-1})\varphi_k(n(t - \frac{k-1}{n})),$$

$$\begin{aligned} \tilde{l}_n(t) &= \frac{1}{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}} \\ &\cdot (x_1 - x_0 + \sum_{m=2}^{k-1} (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)} + \\ &+ (x_k - x_{k-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} n(t - \frac{k-1}{n})) \end{aligned}$$

for $t \in [\frac{k-1}{n}, \frac{k}{n}]$, $(x_1, \dots, x_{n-1}) \in D_n$, $(\varphi_1, \dots, \varphi_n) \in E_\delta^n$.

The function $f = f_n \circ (\tilde{l}_n)^{-1}$ belongs to $\text{Diff}_+^{1,\delta}([0, 1])$, because the left derivation

$$f'((\tilde{l}_n)^{-1}(\frac{k-1}{n} - 0)) = n(x_{k-1} - x_{k-2})\varphi_{k-1}(1).$$

$$\frac{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}}{(x_{k-1} - x_{k-2}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-2}(1)} n} =$$

$$= (x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}) \frac{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)}{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}.$$

is equal to the right derivation

$$f'((\tilde{l}_n)^{-1}(\frac{k-1}{n} + 0)) = n(x_k - x_{k-1})\varphi_k(0).$$

$$\begin{aligned}
& \frac{x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}}{(x_k - x_{k-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} n} = \\
& = (x_1 - x_0 + \sum_{m=2}^n (x_m - x_{m-1}) \frac{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_m(0)}{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{m-1}(1)}) \frac{\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)}{\varphi'_2(0)\varphi'_3(0)\dots\varphi'_{k-1}(0)}.
\end{aligned}$$

Let a subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a). We write

$$L_{\delta,n}(F) = \int_{D_{l_n}} \int_{E_\delta} \dots \int_{E_\delta} F(Q_{l_n}(\bar{x}, \varphi_1, \dots, \varphi_{l_n})) \eta_n(d\bar{x}) \nu(d\varphi_1) \dots \nu(d\varphi_{l_n})$$

for any function $F \in C_b(E_\delta) = C_b(\text{Diff}_+^{1,\delta}([0, 1]))$.

Theorem 3. *If a subgroup G of $\text{Diff}_0^3([0, 1])$ satisfies condition (a) then $\lim_{n \rightarrow \infty} |L_{\delta,n}(F_g) - L_{\delta,n}(F)| = 0$ for any function $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$ and any diffeomorphism $g \in G$.*

Proof. Let $F \in C_b(\text{Diff}_+^{1,\delta}([0, 1]))$, $g \in G$, $C = \sup_{g \in E_\delta} |F(g)|$.

Let $\epsilon \in (0, 1)$.

It follows from Lemma 3 that it exists $\delta_1 \in (0, 1)$ such that

$$\left| \prod_{k=1}^n \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1}) \sqrt{g'(x_k)g'(x_{k-1})}} - 1 \right| \leq \epsilon$$

for any positive integer n and for $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequalities $\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \delta_1$.

Let us take positive ϵ_1 satisfying the inequalities $\epsilon_1 < \frac{1}{8}\epsilon^3$, $\epsilon_1 < \delta_1$, $e^{4c_5 C_g \sqrt[3]{\epsilon_1}} - e^{-4c_4 C_g \sqrt[3]{\epsilon_1}} < \epsilon$.

It follows from Lemma 2 that the inequality is valid $\nu_n(E_\delta^n \setminus X_{\sqrt[3]{\epsilon_1}, g, \bar{x}}) \leq 2\sqrt[3]{\epsilon_1} \leq \epsilon$ for any positive integer n and for any $\bar{x} = (x_1, \dots, x_{n-1}) \in D_n$ satisfying the inequalities $\max_{1 \leq k \leq n} (x_k - x_{k-1}) < \epsilon_1$.

Since the subgroup G satisfies condition (a) we have that

(i) there are a integer $l_n > l_{n-1}$ ($l_0 = 1$) for any natural n and a countably additive Borel measure η_n on D_{l_n} such that $\eta_n(D_{l_n}) = 1$,

(ii) we can find natural $N(\epsilon_1, g)$ such that, for any $n > N(\epsilon_1, g)$, it exists a Borel subset $Z_{n, \epsilon_1, g} \subset D_{l_n}$ that $\eta_n(Z_{n, \epsilon_1, g}) > 1 - \epsilon_1$ and $\max_{1 \leq k \leq l_n} (x_k - x_{k-1}) < \epsilon_1$ for any

$(x_1, x_2, \dots, x_{l_n-1}) \in Z_{n, \epsilon_1, g}$ where $x_0 = 0$, $x_{l_n} = 1$,

(iii) $(1 - \epsilon_1)\eta_n(Y) < \eta_n(gY) < (1 + \epsilon_1)\eta_n(Y)$ for any Borel subset $Y \subset Z_{n, \epsilon_1, g}$ where $gY = \{(g(x_1), g(x_2), \dots, g(x_{l_n-1})) : (x_1, x_2, \dots, x_{l_n-1}) \in Y\}$.

Hence it exists the function $\varrho_n : Z_{n, \epsilon_1, g} \rightarrow \mathbf{R}$ such that $1 - \epsilon_1 \leq \varrho_n(\bar{x}) \leq 1 + \epsilon_1$ for any $\bar{x} \in Z_{n, \epsilon_1, g}$, $\eta_n(gY) = \int_Y \varrho_n(\bar{x}) \eta_n(d\bar{x})$ for any Borel subset $Y \subset Z_{n, \epsilon_1, g}$.

Let $y_k = g(x_k)$, $\bar{y} = (y_1, \dots, y_{l_n-1}) \in D_{l_n}$, $g^{-1}(\bar{y}) = (x_1, \dots, x_{l_n-1}) \in D_{l_n}$. We receive

$$gX_{\sqrt[3]{\epsilon_1}, g, g^{-1}(\bar{y})} = \{(\varphi_1, \dots, \varphi_{l_n}) : (q_1, \dots, q_{l_n}) \in X_{\sqrt[3]{\epsilon_1}, g, \bar{x}},$$

$$Q_{l_n}(y_1, \dots, y_{l_n-1}, \varphi_1, \dots, \varphi_n) = g \circ (Q_{l_n}(x_1, x_2, \dots, x_{l_n-1}, q_1, \dots, q_{l_n}))\}.$$

It is easy to see that $\varphi_k(t) = \frac{g(x_{k-1} + (x_k - x_{k-1})q_k(t)) - g(x_{k-1})}{g(x_k) - g(x_{k-1})}$, because

$$\frac{(y_k - y_{k-1})\varphi'_2(0)\varphi'_3(0)\dots\varphi'_k(0)}{(y_1 - y_0)\varphi'_1(1)\varphi'_2(1)\dots\varphi'_{k-1}(1)} = \frac{(x_k - x_{k-1})q'_2(0)q'_3(0)\dots q'_k(0)}{(x_1 - x_0)q'_1(1)q'_2(1)\dots q'_{k-1}(1)}$$

We have

$$\begin{aligned} & \int_{gZ_{n,\epsilon_1,g}} \nu_{l_n}(gX_{\mathfrak{Z}_{\overline{\epsilon_1},g,g^{-1}(\overline{y})}})\eta_n(d\overline{y}) = \\ &= \int_{Z_{n,\epsilon_1,g}} \left(\int_{X_{\mathfrak{Z}_{\overline{\epsilon_1},g,\overline{x}}}} \exp\left(\sum_{k=1}^{l_n} [(x_k - x_{k-1})\left(\frac{g''(x_{k-1})}{g'(x_{k-1})}q'_k(0) - \frac{g''(x_k)}{g'(x_k)}q'_k(1)\right) + \right. \right. \\ & \left. \left. + (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t))(q'_k(t))^2 dt\right] \nu(dq_1)\dots\nu(dq_{l_n}) \right) \\ & \varrho_n(\overline{x}) \prod_{k=1}^{l_n} \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1})\sqrt{g'(x_k)g'(x_{k-1})}} \eta_n(d\overline{x}) \geq \\ & \geq (1 - \epsilon)^3 \int_{Z_{n,\epsilon_1,g}} \nu_{l_n}(X_{\mathfrak{Z}_{\overline{\epsilon_1},g,\overline{x}}})\eta_n(d\overline{x}) \geq (1 - \epsilon)^5. \end{aligned}$$

Hence,

$$\begin{aligned} & |L_{\delta,n}(F_g) - \int_{gZ_{n,\epsilon_1,g}} \left(\int_{gX_{\mathfrak{Z}_{\overline{\epsilon_1},g,g^{-1}(\overline{y})}}} F_g(Q_{l_n}(\overline{y}, \varphi_1, \dots, \varphi_{l_n})) \right. \\ & \left. \nu(d\varphi_1)\dots\nu(d\varphi_{l_n}) \right) \eta_n(d\overline{y})| \leq C(1 - (1 - \epsilon)^5) \end{aligned}$$

and

$$\begin{aligned} & |L_{\delta,n}(F) - \int_{Z_{n,\epsilon_1,g}} \left(\int_{X_{\mathfrak{Z}_{\overline{\epsilon_1},g,\overline{x}}}} F(Q_{l_n}(\overline{x}, q_1, \dots, q_{l_n})) \right. \\ & \left. \nu(dq_1)\dots\nu(dq_{l_n}) \right) \eta_n(d\overline{x})| \leq C(1 - (1 - \epsilon)^2). \end{aligned}$$

We have

$$\begin{aligned} & \left| \int_{gZ_{n,\epsilon_1,g}} \left(\int_{gX_{\mathfrak{Z}_{\overline{\epsilon_1},g,g^{-1}(\overline{y})}}} F_g(Q_{l_n}(\overline{y}, \varphi_1, \dots, \varphi_{l_n})) \right. \right. \\ & \quad \left. \left. \nu(d\varphi_1)\dots\nu(d\varphi_{l_n}) \right) \eta_n(d\overline{y}) - \int_{Z_{n,\epsilon_1,g}} \left(\int_{X_{\mathfrak{Z}_{\overline{\epsilon_1},g,\overline{x}}}} F(Q_{l_n}(\overline{x}, q_1, \dots, q_{l_n})) \right. \right. \\ & \quad \left. \left. \nu(dq_1)\dots\nu(dq_{l_n}) \right) \eta_n(d\overline{x}) \right| \leq \\ & \leq \int_{Z_{n,\epsilon_1,g}} \left(\int_{X_{\mathfrak{Z}_{\overline{\epsilon_1},g,\overline{x}}}} \left| \exp\left(\sum_{k=1}^n [(x_k - x_{k-1})\left(\frac{g''(x_{k-1})}{g'(x_{k-1})}q'_k(0) - \frac{g''(x_k)}{g'(x_k)}q'_k(1)\right) + \right. \right. \right. \\ & \quad \left. \left. \left. + (x_k - x_{k-1})^2 \int_0^1 S_g(x_{k-1} + (x_k - x_{k-1})q_k(t))(q'_k(t))^2 dt\right] \right) \right. \\ & \quad \left. \varrho_n(\overline{x}) \prod_{k=1}^{l_n} \frac{g(x_k) - g(x_{k-1})}{(x_k - x_{k-1})\sqrt{g'(x_k)g'(x_{k-1})}} - 1 \right| \\ & |F(Q_{l_n}(\overline{x}, q_1, \dots, q_{l_n}))| \nu(dq_1)\dots\nu(dq_n) \eta_n(d\overline{x}) \leq \end{aligned}$$

$$\leq C\epsilon(2 + \epsilon) \int_{Z_n, \epsilon_1, g} \nu_n(X_{\sqrt[3]{\epsilon_1}, g, \bar{x}}) \eta_n(d\bar{x}) \leq C\epsilon(2 + \epsilon),$$

which implies the assertion of Theorem 3.

Define a ultrafilter \mathfrak{S} on the set positive integers such that \mathfrak{S} contains the sets $\{n, n+1, \dots\}$ for any positive integer n . We set $L_\delta(F) = \lim_{\mathfrak{S}} L_{\delta, n}(F)$ for any function $F \in C_b(E_\delta)$.

Note that the limit always exists because $|L_{\delta, n}(F)| \leq \sup_{f \in E_\delta} |F(f)|$.

It is easy to see that $L(e_{1, \delta}) = 1$, $|L_\delta(F)| \leq \sup_{f \in E_\delta} |F(f)|$, and $L(F) \geq 0$ for any nonnegative function $F \in C_b(\text{Diff}_+^{1, \delta}([0, 1]))$. In turn, Theorem 1 follows from Theorem 3.

2. Proof of Theorem 2.

Let $B(G)$ be the linear space of all bounded functions on the group G .

Let positive $\delta < \frac{1}{2}$, let

$$p_\delta(f) = |\ln(f'(0))| + \sup_{t_1, t_2 \in [0, 1]} \frac{|\ln(f'(t_2)) - \ln(f'(t_1))|}{|t_2 - t_1|^\delta}$$

and $r(f) = \inf_{h \in G} p_\delta(h^{-1} \circ f)$ for $f \in \text{Diff}_+^{1, \delta}([0, 1])$, $\theta(t) = 1 - t$ for $t \in [0, 1]$ and $\theta(t) = 0$ for $t > 1$.

For any fixed $f \in \text{Diff}_+^{1, \delta}([0, 1])$, $C > 0$, the set of functions $\{\psi : \psi(t) = \ln(g'(t)), g \in G, p_\delta(g \circ f) < C\}$ contain in a compact subset of the space $C([0, 1])$, therefore it is finite according to condition (a). Hence, we can define the linear mapping $\pi_\delta : B(G) \rightarrow C_b(\text{Diff}_+^{1, \delta}([0, 1]))$ by setting

$$\pi_\delta F(f) = \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f)) F(h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))}.$$

Assign a linear functional $l : B(G) \rightarrow \mathbf{R}$ by setting $l(F) = L_\delta(\pi_\delta F)$.

It is easy to see that

$$|l(F)| = |L_\delta(\pi_\delta F)| \leq \sup_{f \in \text{Diff}_+^{1, \delta}([0, 1])} |\pi_\delta F(f)| \leq \sup_{g \in G} |F(g)|,$$

$l(F) \geq 0$ for any nonnegative function $F \in B(G)$, and $l(e_G) = 1$, where $e_G(g) = 1$ for all $g \in G$.

Denote by $F_g(h) = F(g^{-1} \circ h)$ for $F \in B(G)$, $g, h \in G$.

We have

$$\begin{aligned} \pi_\delta F_g(f) &= \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f)) F(g^{-1} \circ h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ f) - r(f))} = \\ &= \frac{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ g \circ f) - r(g \circ f)) F(h)}{\sum_{h \in G} \theta(p_\delta(h^{-1} \circ g \circ f) - r(g \circ f))} = \pi_\delta F(g \circ f), \end{aligned}$$

hence $l(F_g) = L_\delta(\pi_\delta F_g) = L_\delta(\pi_\delta F) = l(F)$,
which implies the assertion of Theorem 2.

3. Proof of Corollary 2.1.

Let $f_1(t) = \frac{1}{2}t$ for $0 \leq t \leq \frac{1}{2}$, $f_1(t) = t - \frac{1}{4}$ for $\frac{1}{2} \leq t \leq \frac{3}{4}$,
 $f_1(t) = 2t - 1$ for $\frac{3}{4} \leq t \leq 1$ and
 $f_2(t) = t$ for $0 \leq t \leq \frac{1}{2}$, $f_2(t) = \frac{1}{2}t + \frac{1}{4}$ for $\frac{1}{2} \leq t \leq \frac{3}{4}$,
 $f_2(t) = t - \frac{1}{8}$ for $\frac{3}{4} \leq t \leq \frac{7}{8}$, $f_2(t) = 2t - 1$ for $\frac{7}{8} \leq t \leq 1$.

The Thompson's group F is generated by f_1 and f_2 .

Denote by $r_n = 1 - \frac{1}{2^{n+1}}$ for integer $n \geq 0$ and $r_{-k} = \frac{1}{2^k}$ for integer $k \geq 1$. We have $f_1(r_n) = r_{n-1}$ for any integer n .

The group F act on D_n by $f(x_1, \dots, x_{n-1}) = (f(x_1), \dots, f(x_{n-1}))$ for any $f \in F$, $(x_1, \dots, x_{n-1}) \in D_n$.

Let $I_0^k = \{(r_0, r_1)\}$,

$$I_n^k = \{f_2 f_1^{-l_1} f_2 f_1^{l_1-l_2} f_2 \dots f_1^{l_{n-2}-l_{n-1}} f_2 f_1^{l_{n-1}}(r_0, r_1, \dots, r_{n+1}) : 0 \leq l_i \leq \min(k, i)\},$$

$a_{n,k} = |I_n^k|$ for $n \geq 0$, $k \geq 1$.

Lemma 4. $\lim_{n \rightarrow \infty} \frac{a_{n,k}}{4^{n+1} \cos^{2n} \frac{\pi}{k+2}} = (k+2) \sin^2 \frac{\pi}{k+2}$.

Proof. It is easy to see that $a_{0,k} = 1$, $a_{n,1} = 1$, $a_{n+1,k+1} = \sum_{i=0}^n a_{i,k+1} a_{n-i,k}$.

Let $u_k(t) = \sum_{n=0}^{\infty} a_{n,k} t^n$.

We have $u_1(t) = \frac{1}{1-t}$, $u_{k+1}(t) = 1 + t u_{k+1}(t) u_k(t)$ or $u_{k+1}(t) = \frac{1}{1-t u_k(t)}$.

Taking $u_k(t) = \frac{p_{k-1}(t)}{p_k(t)}$, $p_0(t) = 1$, $p_1(t) = 1 - t$ we find

$$u_{k+1}(t) = \frac{1}{1 - t \frac{p_{k-1}(t)}{p_k(t)}} = \frac{p_k(t)}{p_k(t) - t p_{k-1}(t)},$$

$$p_{k+1}(t) = p_k(t) - t p_{k-1}(t).$$

That means

$$p_k(t) = \frac{1}{2^{k+2} \sqrt{1-4t}} [(1 + \sqrt{1-4t})^{k+2} - (1 - \sqrt{1-4t})^{k+2}]$$

$$\text{or } p_k(t) = \prod_{l=1}^{\lfloor \frac{k+1}{2} \rfloor} (1 - 4t \cos^2 \frac{\pi l}{k+2}).$$

Taking $m = \lfloor \frac{k+1}{2} \rfloor$ we find

$$\begin{aligned} u_k(t) &= \frac{4(k+2) \sin^2 \frac{\pi}{k+2}}{1 - 4t \cos^2 \frac{\pi}{k+2}} + \dots + \frac{4(k+2) \sin^2 \frac{\pi m}{k+2}}{1 - 4t \cos^2 \frac{\pi m}{k+2}} = \\ &= \sum_{n=0}^{\infty} 4^{n+1} (k+2) t^n (\sin^2 \frac{\pi}{k+2} \cos^{2n} \frac{\pi}{k+2} + \dots + \sin^2 \frac{\pi m}{k+2} \cos^{2n} \frac{\pi m}{k+2}). \end{aligned}$$

Hence

$$a_{n,k} = 4^{n+1} (k+2) (\sin^2 \frac{\pi}{k+2} \cos^{2n} \frac{\pi}{k+2} + \dots + \sin^2 \frac{\pi m}{k+2} \cos^{2n} \frac{\pi m}{k+2})$$

and $\lim_{n \rightarrow \infty} \frac{a_{n,k}}{4^{n+1} \cos^{2n} \frac{\pi}{k+2}} = (k+2) \sin^2 \frac{\pi}{k+2}$ which implies the assertion of Lemma 4.

For any integer $l \geq 1$, $n_1 \geq 0$, $n_2 \geq 0, \dots, n_l \geq 0$, we write

$$\begin{aligned} Y_{l,n_1,n_2,\dots,n_l} &= \{(r_0, t_{1,1}, t_{1,2}, \dots, t_{1,n_1}, r_1, t_{2,1}, t_{2,2}, \dots, t_{2,n_2}, r_2, \dots, \\ &\quad r_{l-1}, t_{l,1}, t_{l,2}, \dots, t_{l,n_l}, r_l) : (r_{i-1}, t_{i,1}, t_{i,2}, \dots, t_{i,n_i}, r_i) \in f_1^i(I_{n_i}^{l-i}), \quad 0 \leq i \leq l\}, \\ Y^{l,n} &= \bigcup_{n_1+\dots+n_l=n, n_1 \geq 0, \dots, n_l \geq 0} Y_{l,n_1,\dots,n_l}, \\ Y_0^{l,n} &= \bigcup_{n_2+\dots+n_l=n, n_2 \geq 0, \dots, n_l \geq 0} Y_{l,0,n_2,\dots,n_l}. \end{aligned}$$

It is easy to see that $f_2(Y^{l+1,n}) = Y^{l,n+1} \setminus Y_0^{l,n+1}$.

Introduce the mapping $\kappa_n : D_n \rightarrow D_{2n}$ by setting

$$\kappa_n(x_1, x_2, \dots, x_{n-1}) = \left(\frac{x_1}{2}, x_1, \frac{x_1+x_2}{2}, x_2, \frac{x_2+x_3}{2}, \dots, \frac{x_{n-2}+x_{n-1}}{2}, x_{n-1}, \frac{x_{n-1}+1}{2}\right).$$

Denote by $X^{0,l,n} = \bigcup_{i=0}^l \bigcup_{j=0}^{l-1} f_1^j(Y^{2l-i,n+i})$,

$$X^{m,l,n} = \kappa_{2^{m-1}(n+2l+2)}(\kappa_{2^{m-2}(n+2l+2)}(\dots \kappa_{2(n+2l+2)}(\kappa_{n+2l+2}(X^{0,l,n}))\dots)).$$

We have $X^{m,l,n} = \kappa_{2^{m-1}(n+2l+2)}(X^{m-1,l,n})$ and $f_1(\kappa_{2^{m-1}(n+2l+2)}(\bar{x})) = \kappa_{2^{m-1}(n+2l+2)}(f_1(\bar{x}))$, $f_2(\kappa_{2^{m-1}(n+2l+2)}(\bar{x})) = \kappa_{2^{m-1}(n+2l+2)}(f_2(\bar{x}))$ for any $\bar{x} \in X^{m-1,l,n}$.

Also, if $(t_1, t_2, \dots, t_{2^m(n+2l+2)})$ belongs to $X^{m,l,n}$ then

$$\{t_1, t_2, \dots, t_{2^m(n+2l+2)}\} \supset \left\{\frac{1}{2^m}, \frac{2}{2^m}, \frac{3}{2^m}, \dots, \frac{2^m-1}{2^m}\right\}$$

for any $m \geq 1$.

Lemma 5. *For any positive ε , there are positive integer l, n such that*

$$\frac{|f_1(X^{m,l,n}) \cap X^{m,l,n}|}{|X^{m,l,n}|} > 1 - \varepsilon, \quad \frac{|f_2(X^{m,l,n}) \cap X^{m,l,n}|}{|X^{m,l,n}|} > 1 - \varepsilon \text{ for any } m \geq 0.$$

Proof. It is sufficient to prove for $m = 0$. Take integer $l > \frac{4}{\varepsilon}$.

It follows from Lemma 4 that it exists such integer n that $\frac{|Y_0^{2l-i,n+i}|}{|Y^{2l-i,n+i}|} < \frac{1}{l^3}$ for any $0 \leq i \leq l$.

As $f_2(Y^{2l-i,n+i}) = Y^{l-i-1,n+i+1} \setminus Y_0^{l-i-1,n+i+1}$ we have

$$(1 - \frac{1}{l^2})|Y^{l,n+l}| \leq |Y^{2l,n}| \leq |Y^{2l-1,n+2}| \leq \dots \leq |Y^{l,n+l}|$$

and $\frac{|Y^{l,n+l}|}{|\bigcup_{i=0}^l Y^{2l-i,n+i}|} < \frac{1}{l}$.

$$\text{Hence, } \frac{|f_2(X^{0,l,n}) \cap X^{0,l,n}|}{|X^{0,l,n}|} > 1 - \frac{|Y^{l,n+l}|}{|\bigcup_{i=0}^l Y^{2l-i,n+i}|} > 1 - \frac{1}{l} > 1 - \varepsilon.$$

$$\text{As } f_1\left(\bigcup_{j=0}^{l-1} f_1^j(Y^{2l-i,n+i})\right) \cap \bigcup_{j=0}^{l-1} f_1^j(Y^{2l-i,n+i}) = \bigcup_{j=1}^{l-1} f_1^j(Y^{2l-i,n+i})$$

we find $\frac{|f_1(X^{0,l,n}) \cap X^{0,l,n}|}{|X^{0,l,n}|} = 1 - \frac{1}{l} > 1 - \varepsilon$ which implies the assertion of Lemma 5.

Take a infinite differential function $\psi : \mathbf{R} \rightarrow \mathbf{R}$ such that $\psi(t+1) = \psi(t) + 2$, $0 < \psi'(t) \leq 3$ for any $t \in \mathbf{R}$, $\psi'(t) = 3$ for any $t \in [\frac{1}{4}, \frac{3}{4}]$, $\psi(0) = 0$, $\psi(\frac{1}{4}) = \frac{1}{4}$, $\psi'(0) = 1, \psi^{(n)}(0) = 0$ for any $n \geq 2$.

For any dyadic rational $r = \frac{k}{2^p} \in (0, 1)$, denote $x_r = \psi^{-p}(k)$, $x'_r = \psi^{-p}(k - \frac{1}{4})$, $x''_r = \psi^{-p}(k + \frac{1}{4})$, $\phi_r(t) = \psi^{-p}(k + t)$.

Let $g_1(t) = \psi^{-1}(t)$ for $0 \leq t \leq x_{\frac{1}{2}} = \frac{1}{2}$,
 $g_1(t) = \psi^{-2}(\psi^2(t) - 1)$ for $x_{\frac{1}{2}} \leq t \leq x_{\frac{3}{4}}$,
 $g_1(t) = \psi(t) - 1$ for $x_{\frac{3}{4}} \leq t \leq 1$ and
 $g_2(t) = t$ for $0 \leq t \leq x_{\frac{1}{2}}$,
 $g_2(t) = \psi^{-2}(\psi(t) + 1)$ for $x_{\frac{1}{2}} \leq t \leq x_{\frac{3}{4}}$,
 $g_2(t) = \psi^{-3}(\psi^3(t) - 1)$ for $x_{\frac{3}{4}} \leq t \leq x_{\frac{7}{8}}$,
 $g_2(t) = \psi(t) - 1$ for $x_{\frac{7}{8}} \leq t \leq 1$.

In [2] È.Ghys and V.Sergiescu proved that the Thompson's group F is isomorphic to a discrete subgroup G of $\text{Diff}_0^3([0, 1])$ which is generated by $\{g_1, g_2\}$ and satisfies condition (b).

Lemma 6. *For any dyadic rational $r \in (0, 1)$, there are positive integer $\alpha_1, \alpha_2, \beta_1, \beta_2$ such that $|\alpha_1| \leq 1, |\alpha_2| \leq 1, |\beta_1| \leq 1, |\beta_2| \leq 1$,
 $g_1(\phi_r(t)) = \phi_{f_1(r)}(\psi^{\alpha_1}(t)), g_2(\phi_r(t)) = \phi_{f_2(r)}(\psi^{\beta_1}(t)), g_1(\phi_r(-t)) = \phi_{f_1(r)}(\psi^{\alpha_2}(-t)),$
 $g_2(\phi_r(-t)) = \phi_{f_2(r)}(\psi^{\beta_2}(-t))$ for any $t \in [0, \frac{1}{4}]$.*

Proof. Let $t \in [0, \frac{1}{4}]$.

If $r = \frac{1}{2}$ we have $f_1(r) = \frac{1}{4}, f_2(r) = \frac{1}{2}$,

$$\begin{aligned} g_1(\phi_r(t)) &= \psi^{-2}(\psi^2(\psi^{-1}(t+1)) - 1) = \psi^{-2}(\psi(t) + 1) = \phi_{f_1(r)}(\psi(t)), \\ g_2(\phi_r(t)) &= \psi^{-2}(\psi((\psi^{-1}(t+1))) + 1) = \psi^{-1}((\psi^{-1}(t) + 1) = \phi_{f_2(r)}(\psi^{-1}(t)), \\ g_1(\phi_r(-t)) &= \psi^{-1}(\psi^{-1}(-t+1)) = \psi^{-2}(-t+1) = \phi_{f_1(r)}(-t), \\ g_2(\phi_r(-t)) &= \psi^{-1}(-t+1) = \phi_{f_2(r)}(-t). \end{aligned}$$

Hence $\alpha_1 = 1, \alpha_2 = 0, \beta_1 = -1, \beta_2 = 0$.

If $r = \frac{3}{4}$ we have $f_1(r) = \frac{1}{2}, f_2(r) = \frac{5}{8}$,

$$\begin{aligned} g_1(\phi_r(t)) &= \psi(\psi^{-2}(t+3) - 1) = \psi^{-1}(t+1) = \phi_{f_1(r)}(t), \\ g_2(\phi_r(t)) &= \psi^{-3}(\psi^3(\psi^{-2}(t+3)) - 1) = \psi^{-3}((\psi(t) + 5) = \phi_{f_2(r)}(\psi(t)), \\ g_1(\phi_r(-t)) &= \psi^{-2}(\psi^2(\psi^{-2}(-t+3)) - 1) = \psi^{-1}(\psi^{-1}(-t) + 1) = \phi_{f_1(r)}(\psi^{-1}(-t)), \\ g_2(\phi_r(-t)) &= \psi^{-2}(\psi(\psi^{-2}(-t+3)) + 1) = \psi^{-3}(-t+5) = \phi_{f_2(r)}(-t). \end{aligned}$$

Hence $\alpha_1 = 0, \alpha_2 = -1, \beta_1 = 1, \beta_2 = 0$.

If $r = \frac{7}{8}$ we have $f_2(r) = \frac{3}{4}$,

$$\begin{aligned} g_2(\phi_r(t)) &= \psi(\psi^{-3}(t+7)) - 1 = \psi^{-2}(t+3) = \phi_{f_2(r)}(t), \\ g_2(\phi_r(-t)) &= \psi^{-3}(\psi^3(\psi^{-3}(-t+7)) - 1) = \psi^{-2}(\psi^{-1}(-t) + 3) = \phi_{f_2(r)}(-t). \end{aligned}$$

Hence $\beta_1 = 0, \beta_2 = -1$.

If $0 < r = \frac{k}{2^p} < \frac{1}{2}$ we have $f_1(r) = \frac{k}{2^{p+1}}, f_2(r) = \frac{k}{2^p}$,

$$\begin{aligned} g_1(\phi_r(\pm t)) &= \psi^{-1}(\psi^{-p}(\pm t + k)) = \psi^{-p-1}(\pm t + k) = \phi_{f_1(r)}(\pm t), \\ g_2(\phi_r(-t)) &= \psi^{-p}(\pm t + k) = \phi_{f_2(r)}(\pm t). \end{aligned}$$

Hence $\alpha_1 = \alpha_2 = 0, \beta_1 = \beta_2 = 0$.

If $\frac{1}{2} < r = \frac{k}{2^p} < \frac{3}{4}$ we have $f_1(r) = \frac{k-2^{p-2}}{2^p}, f_2(r) = \frac{k+2^{p-1}}{2^{p+1}}$,

$$\begin{aligned} g_1(\phi_r(\pm t)) &= \psi^{-2}(\psi^2(\psi^{-p}(\pm t + k)) - 1) = \psi^{-p}(\pm t + k - 2^{p-2}) = \phi_{f_1(r)}(\pm t), \\ g_2(\phi_r(\pm t)) &= \psi^{-2}(\psi(\psi^{-p}(\pm t + k)) + 1) = \psi^{-p-1}(\pm t + k + 2^{p-1}) = \phi_{f_2(r)}(\pm t). \end{aligned}$$

Hence $\alpha_1 = \alpha_2 = 0, \beta_1 = \beta_2 = 0$.

If $\frac{3}{4} < r = \frac{k}{2^p} < 1$ we have $f_1(r) = \frac{k-2^{p-1}}{2^{p-1}}$,

$$g_1(\phi_r(\pm t)) = \psi(\psi^{-p}(\pm t + k)) - 1 = \psi^{-p+1}(\pm t + k - 2^{p-1}) = \phi_{f_1(r)}(\pm t).$$

Hence $\alpha_1 = \alpha_2 = 0$.

If $\frac{3}{4} < r = \frac{k}{2^p} < \frac{7}{8}$ we have $f_2(r) = \frac{k-2^{p-3}}{2^p}$,

$$g_2(\phi_r(\pm t)) = \psi^{-3}(\psi^3(\psi^{-p}(\pm t + k)) - 1) = \psi^{-p}(\pm t + k - 2^{p-3}) = \phi_{f_2(r)}(\pm t).$$

Hence $\beta_1 = \beta_2 = 0$.

If $\frac{7}{8} < r = \frac{k}{2^p} < 1$ we have $f_2(r) = \frac{k-2^{p-1}}{2^{p-1}}$,

$$g_2(\phi_r(\pm t)) = \psi(\psi^{-p}(\pm t + k)) - 1 = \psi^{-p+1}(\pm t + k - 2^{p-1}) = \phi_{f_2(r)}(\pm t).$$

Hence $\beta_1 = \beta_2 = 0$.

Thus, we prove Lemma 6.

Lemma 7. For any positive ε , there are positive integer N and a finite subset $Z \subset D_N$ such that $\frac{|g_1(Z) \cap Z|}{|Z|} > 1 - \varepsilon$, $\frac{|g_2(Z) \cap Z|}{|Z|} > 1 - \varepsilon$,

$\max_{1 \leq k \leq N} (x_k - x_{k-1}) < \varepsilon$ for any $(x_1, x_2, \dots, x_{N-1}) \in Z$ where $x_0 = 0$, $x_N = 1$.

Proof. Let $\varepsilon \in (0, 1)$.

As $\lim_{m \rightarrow \infty} \sum_{l=1}^{2^m-1} (x''_{\frac{l}{2^m}} - x'_{\frac{l}{2^m}}) = \frac{1}{2}$ it exists such $m \geq 1$ that $\max_{1 \leq l \leq 2^m} (x'_{\frac{l}{2^m}} - x''_{\frac{l}{2^m}}) < \varepsilon$,

where $x''_0 = \frac{1}{4}$, $x'_1 = \frac{3}{4}$.

By Lemma 5 we find positive integer l, n such that

$$\frac{|f_1(X^{m,l,n}) \cap X^{m,l,n}|}{|X^{m,l,n}|} > 1 - \frac{1}{4}\varepsilon, \quad \frac{|f_2(X^{m,l,n}) \cap X^{m,l,n}|}{|X^{m,l,n}|} > 1 - \frac{1}{4}\varepsilon.$$

Let $k = 2^m(n+2l+2)$, $V_{\bar{t}} = \{t_1, t_2, \dots, t_{k-1}\}$ for any $\bar{t} = (t_1, t_2, \dots, t_{k-1}) \in X^{m,l,n}$, and $W = \bigcup_{\bar{t} \in X^{m,l,n}} V_{\bar{t}}$.

Take integer $J > \frac{16(k+1)}{\varepsilon}$. Let

$$C = \max_{0 \leq j \leq J} \left(\max_{-\frac{1}{4} \leq x \leq \frac{1}{4}} \left(\max_{r \in W} |(\phi_r(\psi^j(x)))'| + |(\psi^j(x))'| \right) \right).$$

Take integer $p > \frac{C+1}{\varepsilon}$. Let $N = k(2p+1)$,

$$Z = \{(\psi^{j_1}(\frac{1}{4p}), \psi^{j_1}(\frac{2}{4p}), \dots, \psi^{j_1}(\frac{p-1}{4p}), \frac{1}{4}, x'_{t_1}, \phi_{t_1}(\psi^{j_2}(-\frac{p-1}{4p})), \phi_{t_1}(\psi^{j_2}(-\frac{p-2}{4p})), \dots,$$

$$\phi_{t_1}(\psi^{j_2}(-\frac{1}{4p})), x_{t_1}, \phi_{t_1}(\psi^{j_3}(\frac{1}{4p})), \phi_{t_1}(\psi^{j_3}(\frac{2}{4p})), \dots, \phi_{t_1}(\psi^{j_3}(\frac{p-1}{4p})), x''_{t_1},$$

$$x'_{t_2}, \phi_{t_2}(\psi^{j_4}(-\frac{p-1}{4p})), \phi_{t_2}(\psi^{j_4}(-\frac{p-2}{4p})), \dots, \phi_{t_2}(\psi^{j_4}(-\frac{1}{4p})),$$

$$x_{t_2}, \phi_{t_2}(\psi^{j_5}(\frac{1}{4p})), \phi_{t_2}(\psi^{j_5}(\frac{2}{4p})), \dots, \phi_{t_2}(\psi^{j_5}(\frac{p-1}{4p})), x''_{t_2}, \dots,$$

$$x'_{t_{k-1}}, \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{p-1}{4p})), \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{p-2}{4p})), \dots, \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{1}{4p})),$$

$$x_{t_{k-1}}, \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{1}{4p})), \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{2}{4p})), \dots, \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{p-1}{4p})), x''_{t_{k-1}}, \dots,$$

$$\frac{3}{4}, 1 - \psi^{j_{2k}}(-\frac{p-1}{4p}), 1 - \psi^{j_{2k}}(-\frac{p-2}{4p}), \dots, 1 - \psi^{j_{2k}}(-\frac{1}{4p}) : 0 \leq j_1 \leq J, 0 \leq j_2 \leq J,$$

$$0 \leq j_3 \leq J, 0 \leq j_4 \leq J, 0 \leq j_5 \leq J, \dots, 0 \leq j_{2k-2} \leq J, 0 \leq j_{2k-1} \leq J,$$

$$0 \leq j_{2k} \leq J, (t_1, t_2, \dots, t_{k-1}) \in X^{m,l,n}\},$$

and

$$\begin{aligned}
Z_i = & \{(\psi^{j_1}(\frac{1}{4p}), \psi^{j_1}(\frac{2}{4p}), \dots, \psi^{j_1}(\frac{p-1}{4p}), \frac{1}{4}, x'_{t_1}, \phi_{t_1}(\psi^{j_2}(-\frac{p-1}{4p})), \phi_{t_1}(\psi^{j_2}(-\frac{p-2}{4p})), \dots, \\
& \phi_{t_1}(\psi^{j_2}(-\frac{1}{4p})), x_{t_1}, \phi_{t_1}(\psi^{j_3}(\frac{1}{4p})), \phi_{t_1}(\psi^{j_3}(\frac{2}{4p})), \dots, \phi_{t_1}(\psi^{j_3}(\frac{p-1}{4p})), x''_{t_1}, \\
& x'_{t_2}, \phi_{t_2}(\psi^{j_4}(-\frac{p-1}{4p})), \phi_{t_2}(\psi^{j_4}(-\frac{p-2}{4p})), \dots, \phi_{t_2}(\psi^{j_4}(-\frac{1}{4p})), \\
& x_{t_2}, \phi_{t_2}(\psi^{j_5}(\frac{1}{4p})), \phi_{t_2}(\psi^{j_5}(\frac{2}{4p})), \dots, \phi_{t_2}(\psi^{j_5}(\frac{p-1}{4p})), x''_{t_2}, \dots, \\
& x'_{t_{k-1}}, \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{p-1}{4p})), \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{p-2}{4p})), \dots, \phi_{t_{k-1}}(\psi^{j_{2k-2}}(-\frac{1}{4p})), \\
& x_{t_{k-1}}, \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{1}{4p})), \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{2}{4p})), \dots, \phi_{t_{k-1}}(\psi^{j_{2k-1}}(\frac{p-1}{4p})), x''_{t_{k-1}}, \dots, \\
& \frac{3}{4}, 1-\psi^{j_{2k}}(-\frac{p-1}{4p}), 1-\psi^{j_{2k}}(-\frac{p-2}{4p}), \dots, 1-\psi^{j_{2k}}(-\frac{1}{4p}) : 1 \leq j_1 \leq J-1, 1 \leq j_2 \leq J-1, \\
& 1 \leq j_3 \leq J-1, 1 \leq j_4 \leq J-1, 1 \leq j_5 \leq J-1, \dots, 1 \leq j_{2k-2} \leq J-1, 1 \leq j_{2k-1} \leq J-1, \\
& 0 \leq j_{2k} \leq J, (t_1, t_2, \dots, t_{k-1}) \in f_i(X^{m,l,n}) \bigcap X^{m,l,n}\}
\end{aligned}$$

where $i = 1, 2$.

By Lemma 6 we find $g_1(Z_1) \subset Z$ and $g_2(Z_2) \subset Z$. Hence

$$\begin{aligned}
\frac{|g_1(Z) \cap Z|}{|Z|} & \leq \frac{|Z_1|}{|Z|} = \frac{(J-1)^{(2k)} |f_1(X^{m,l,n}) \cap X^{m,l,n}|}{(J+1)^{(2k)} |X^{m,l,n}|} > \\
& > (1 - \frac{4k}{J+1})(1 - \frac{1}{4}\varepsilon) > (1 - \frac{1}{4}\varepsilon)^2 > 1 - \varepsilon, \\
\frac{|g_2(Z) \cap Z|}{|Z|} & \leq \frac{|Z_2|}{|Z|} = \frac{(J-1)^{(2k)} |f_2(X^{m,l,n}) \cap X^{m,l,n}|}{(J+1)^{(2k)} |X^{m,l,n}|} > 1 - \varepsilon.
\end{aligned}$$

We have $\phi_r(\psi^j(\frac{i}{4p})) - \phi_r(\psi^j(\frac{i-1}{4p})) \leq C \frac{1}{4p} < \varepsilon$, $\psi^j(\frac{i}{4p}) - \psi^j(\frac{i-1}{4p}) \leq C \frac{1}{4p} < \varepsilon$ for any $r \in W$, $1 \leq i \leq p$, $1 \leq j \leq J$ that means $\max_{1 \leq k' \leq N} (x_{k'} - x_{k'-1}) < \varepsilon$ for any $(x_1, x_2, \dots, x_{N-1}) \in Z$.

Thus, we prove Lemma 7.

In turn, Corollary 2.1 follows from Theorem 2 and Lemma 7.

Acknowledgements. The author acknowledge I.K. Babenko, M.Brin, K.Brown, R. Grigorchuk, V.S. Guba, P. de la Harpe, J.Moore, M.Sapir, V.Sergiescu, O.G. Smolyanov, A.I. Stern for the discussions and interest about this work.

References

1. Cannon J.W., Floyd W.J., Parry W.R. "Introductory notes on Richard Thompson's groups Enseign. Math., vol 42, issue 2 (1996), pages 215–256.
2. Ghys È., Sergiescu V. "Sur un groupe remarquable de difféomorphismes du cercle Comment.Math.Helvetici 62 (1987) 185–239.
3. Shavgulidze E.T. "Some Properties of Quasi-Invariant Measures on Groups of Diffeomorphisms of the Circle Russ. J. Math. Physics, vol 7, issue 4 (2000), pages 464–472.
4. Hui-Hsiuhg Kuo "Gaussian measures in Banach space Lecture Notes in Mathematics 463, Springer-Verlag 1975.